

Primary Mailing Address: 10820 State Road 54, Ste. 201 Trinity, FL, 34655-1783 Gastroenterology Associates of West Florida

Digestive Diseases and Nutrition

Diplomates American Board of Gastroenterology Diplomates American Board of Internal Medicine Curtis Freedland, D.O. Gastroenterology & Nutrition Joseph Staffetti, M.D. Gastroenterology Dilip Ghanekar, M.D. Gastroenterology

(727) 846-7031 Fax (727) 846-7132

Preparing for Colonoscopy or Flexible Sigmoidoscopy using the Two Day MiraLax Prep

You are scheduled for a colonoscopy or a flexible sigmoidoscopy on

> The Surgery Center will call you one day prior to the procedure with the time.

Please be courteous to others and call with cancellations at least 72 hours prior to the procedure. Make appropriate arrangements for copay and deductibles with the surgery center.

- > WE WILL NOT PERFORM THE EXAM ON PATIENTS WHO COME ALONE
 - You <u>will not be able to drive home after the test</u> if you receive sedation. You <u>must be</u> <u>accompanied</u> by someone who can see you home after the exam.
- You need to <u>HOLD</u> Coumadin, Plavix, Aspirin, Fish Oil, Vitamin E, multivitamins, Arthritis Pills, and Ibuprophen unless instructed otherwise for five (5) days prior to the exam. You may take Tylenol and/or Darvocet for pain.

Items to be purchased prior to the exam (all available over the counter):

One bottle of Citrate of Magnesia

- > Dulcolax Laxatives, 8 tablets
- MiraLax 238 /8.3 oz grams
- 64 Oz of Gatorade or PowerAde, Crystal Light, or vitamin water (Not Red or carbonated)

Two days prior to the exam:

- 1. Take clear liquids only ALL DAY such as soda, broth, apple juice, tea, **BLACK** coffee, or ice pops.
- 2. Take one (1) bottle of citrate of magnesia at about 3 p.m.
- 3. Take 2 Dulcolax laxatives by mouth at 7 p.m.

One day prior to the exam:

- 1. Take clear liquids ALL DAY: Soda, broth, apple juice, tea & coffee BLACK (NO Creamer sugar is okay) and ice pops NOTHING THE COLOR RED AND NOTHING THAT CONTAINS ALCOHOL.
- 2. Put the 64 oz. of Gatorade or PowerAde (No Red or Carbonated Beverages) into a large pitcher at room temperature add the entire contents of MiraLax to the pitcher. Mix and refrigerate.
- 3. Continue to drink plenty of liquids throughout the day and evening.
- 4. At 4 p.m. Drink 8 oz. of Gatorade or PowerAde mixed with MiraLax every 15 minutes until the entire 64 oz. is consumed.
- 5. At 7:00 p.m. take 2 Dulcolax tablets by mouth. If not running clear, take 2 Dulcolax tablets at midnight (clear is no solid pieces, may be Yellow, Green, liquid or mucus, but no solids).
- 6. Nothing by mouth after midnight.

Day of the exam:

1. Take only blood pressure, heart, and lung medications as discussed with your doctor with a sip of water. Do not take diabetes pills or insulin (See Diabetic instructions).

2. Do not wear makeup or nail polish.

**Buy some flushable baby wipes and put in refrigerator, use to help soothe the rectum. **

Doctor Doing Procedure:

Patient Signature:

Date:

Doctors are available 24 hours day or night. Please call if you have problems with prep, questions about medications, or any other concerns: Call (727).846.7031